

Dr. MARIA JOSE PINZON GARCIA
RED COLOMBIANA DE MEDICINA GENETICA SAS - PREGEN
BOGOTA
CARRERA 15 A # 106 - 42
11001 BOGOTA
Colombia

Date of Report 29.05.2026
Sample Received 26.05.2026
Date of Sampling 14.05.2026
LAB-ID 262023695

Medical Report

| | | | |
|---------------|--------------------------------|-----------|----------|
| Patient name | GAMBOA ARRIETA VICTORIA | Sample-ID | A0341495 |
| Date of Birth | 21.03.2026 | Gender | F |

Indication: Newborn Screening

Method(s): Immunoassay, Tandem mass spectrometry from Dried Blood Spot. qPCR from Dried Blood Spot.

Results:

| Parameter | Value | Unit | Reference |
|--|----------|--------|-----------|
| Birth weight (g) | 3600 | g | - |
| 17-hydroxyprogesterone (17OHP) | <5.0 | nmol/L | < 90.0 |
| Thyroid-stimulating hormone (TSH) | <0.7 | µU/mL | < 15.0 |
| Biotinidase | 269.1 | U | > 51.0 |
| Galactose-1-P-uridyltransferase (GALT) | 6.2 | U/g Hb | > 2.5 |
| Immunoreactive trypsinogen (IRT) | <15 | ng/mL | < 65.0 |
| Phenylalanine | 40.9 | µmol/L | < 150.0 |
| Amino acid profile | negative | | - |
| Acylcarnitine profile | negative | | - |

Interpretation: NEGATIVE RESULT

| | |
|---------------|--------------------------------|
| Patient name | GAMBOA ARRIETA VICTORIA |
| Date of Birth | 21.03.2026 |

| | |
|-----------|----------|
| Sample-ID | A0341495 |
| Gender | F |

Results:

Amino Acids

| Parameter | Value | Unit | Reference |
|--|-------|--------|--------------|
| Phenylalanine (Phe) | 40.9 | µmol/L | < 150.0 |
| Phenylalanine / Tyrosine ratio (Phe/Tyr) | 0.47 | µmol/L | < 2.20 |
| Tyrosine (Tyr) | 87.0 | µmol/L | < 200.0 |
| Leucine (Leu) | 122.4 | µmol/L | < 270.0 |
| Valine (Val) | 45.1 | µmol/L | < 200.0 |
| Methionine (MET) | 21.9 | µmol/L | < 78.0 |
| Methionine / Phenylalanine (Met/Phe) | 0.54 | µmol/L | < 1.60 |
| Citrulline (Cit) | 10.7 | µmol/L | < 50.0 |
| Ornithine (Orn) | 100.4 | µmol/L | < 250.0 |
| Ornithine / Citrulline ratio (Orn/Cit) | 9.38 | µmol/L | 1.50 - 20.00 |
| Proline (Pro) | 132.2 | µmol/L | < 350.0 |
| Alanine (Ala) | 126.1 | µmol/L | < 750.0 |
| Arginine (Arg) | 33.9 | µmol/L | < 100.0 |
| Aspartic acid (Asp) | 135.0 | µmol/L | < 100.0 |
| Glutamic acid (Glu) | 212.9 | µmol/L | < 600.0 |
| Glycamine (Gly) | 116.4 | µmol/L | < 700.0 |

Acylcarnitines

| | | | |
|--|-------|--------|---------------|
| Free carnitine (C0) | 31.01 | µmol/L | 6.00 - 100.00 |
| acetylcarnitine (C2) | 12.15 | µmol/L | 1.34 - 48.81 |
| propionylcarnitine (C3) | 1.11 | µmol/L | 0.13 - 6.60 |
| butyryl-/isobutyrylcarnitine (C4) | 0.12 | µmol/L | 0.03 - 0.90 |
| isovaleryl-/2-methylbutyrylcarnitine(C5) | 0.15 | µmol/L | 0.02 - 2.00 |
| tiglylcarnitine (C5:1) | 0.02 | µmol/L | < 0.20 |
| hydroxyvalerylcarnitine (C5OH) | 0.41 | µmol/L | 0.02 - 0.57 |
| glutarylacarnitine (C5DC) | 0.04 | µmol/L | < 0.30 |
| hexanoylcarnitine (C6) | 0.03 | µmol/L | 0.01 - 0.13 |
| octanoylcarnitine (C8) | 0.03 | µmol/L | 0.01 - 0.30 |
| decanoylcarnitine (C10) | 0.04 | µmol/L | 0.01 - 0.36 |
| decenoylcarnitine (C10:1) | 0.09 | µmol/L | < 0.30 |
| dodecanoylcarnitine (C12) | 0.05 | µmol/L | 0.10 - 0.60 |
| myristoylcarnitine (C14) | 0.09 | µmol/L | 0.01 - 0.57 |
| tetradecenoylcarnitine (C14:1) | 0.09 | µmol/L | 0.10 - 0.38 |
| palmitoylcarnitine (C16) | 0.54 | µmol/L | 0.62 - 7.81 |
| 3-hydroxypalmitoylcarnitine (C16OH) | 0.02 | µmol/L | < 0.10 |
| stearoylcarnitine (C18) | 0.26 | µmol/L | 0.30 - 2.40 |
| oleylcarnitine (C18:1) | 2.32 | µmol/L | 0.06 - 3.86 |
| 3-hydroxystearoylcarnitine (C18OH) | 0.01 | µmol/L | < 0.09 |
| malonylcarnitine (C3DC) | 0.03 | µmol/L | < 0.50 |

Amino acid levels are indicators of phenylketonuria, tyrosinemia, MSUD, hydroxyprolinuria, hypermethioninemia (homocystinuria), citrullinemia, argininosuccinate aziduria, hyperargininemia, and hyperprolinemia. Acylcarnitine levels are indicators of carnitine uptake disorders, CPT-I deficiency, CPT-II deficiency, CAT deficiency, propionaciduria, methylmalonic aciduria, malonic aciduria, SCAD deficiency/ethylmalonic aciduria, isovaleric aciduria, HMG-CoA lyase deficiency, 3-methylcrotonyl-CoA carboxylase deficiency, methylglutaconiduria, MCAD deficiency, VLCAD deficiency, LCHAD deficiency, glutaraziduria I, multiple acyl-CoA dehydrogenase deficiency (MAD deficiency/glutaraziduria II), and Beta-ketothiolase deficiency.

Please note: Inconspicuous negative biochemical results cannot exclude any inborn error of metabolism or endocrine disorder with certainty in newborns. We recommend any follow-up or genetic testing if any clinical symptoms are present.

Authorized By: Assoc.-Prof. Dr. Andrea-Romana KASPER, MD, PhD
[Specialist for Pediatrics, Neonatology and Nutrition]

Report was electronically signed and approved.

Contact Details

Assoc.-Prof. Dr. Andrea-Romana KASPER, MD, PhD
E-Mail: info@archimedlife.com

ARCHIMEDlife GmbH
International Medical Laboratory+
Leberstrasse 20/2 | 1110 Vienna, Austria
www.archimedlife.com